

Criteria 2022



Saving wild ocean places, for us and future generations

Table of Contents

	Introduction to Blue Park Awards			2
Ш	The Evaluation Process			4
Ш	The Award Criteria			7
	1	1.1 1.2	Biodiversity Value Implementation	10 10 11
	-	2.1 2.2	Regulations Design, Management & Capacity	13 18
	3	Syst (3.1 3.2	em Priorities Ecosystem Representation Ecological Spatial Connectivity	19 19 19
IV	The Scient	tific Ba	sis for Blue Park Criteria	20
	1	Eligil	bility Criteria	21
		1.1 1.2	Biodiversity Value Implementation	21 22
	2	Awar	d Status Criteria	23
		2.1 2.2	Regulations Design, Management & Capacity	23 24
	3	Syst	em Priorities	25
		3.1 3.2	Ecosystem Representation Ecological Spatial Connectivity	26 26
V	Acknowledgements			27
VI	References 29			29



Introduction to Blue Park Awards

Introduction to Blue Park Awards

Blue Park Award Criteria Marine Conservation Institute aims to safeguard marine biodiversity by assembling a strategic network of effective marine protected areas (MPAs) that includes replicate representation of all marine habitats in each biogeographic region and supports marine population persistence.

To make this vision a reality, we must improve the quality of MPAs and accelerate the implementation of MPAs. Despite the implementation of more than 11,000 MPAs around the world.

less than 3% of the ocean is strongly protected.

(MPAtlas.org)


These efforts are not keeping pace with the growing threats to marine life.

Blue Park Awards incentivize the implementation of more effective MPAs by awarding those that meet science-based standards for effectiveness. Governments and NGOs welcome Blue Park Awards because they attract tourists, local support and investors, and they bring a sense of pride to political leaders, managers, and communities. Blue Park Awards support MPA efforts around the world, complementing the work of communities, leaders, conservation groups and governments to implement MPAs and fulfill international MPA commitments.



The Evaluation Process

The Evaluation Process



Blue Park Award Criteria Any MPA that meets the science-based Blue Park standards defined by the following criteria qualifies for an award and inclusion in the Blue Park network. Becoming a Blue Park is a four-step process. First, MPAs are nominated through an online nomination platform that elicits information and documentation needed for the evaluation. Second, program staff complete an evaluation report for the nominee based on the criteria. During this step, program staff reach out to MPA managers to invite their contributions to the evaluation report. The evaluation report includes recommendations to improve the efficacy of the MPA with respect to safeguarding and recovering marine biodiversity. Evaluation reports are published on the Blue Parks web page for 30 days, during which time, those with experience in the nominated MPA are encouraged to submit comments as well as additional documentation or evidence to correct and refine the report. Third, the evaluation report and the collected comments are forwarded to Science Council members with expertise in the nominee's region and ecosystems. Science Council members determine the nominee's award status.

Nominees earning a Blue Park Award are publicly recognized and celebrated by Blue Park partners. All Blue Parks are subject to a review audit every five years. An earlier audit may be triggered when significant changes to a Blue Park are brought to the attention of program staff.

NOMINATION



MPA nominated through Blue Parks web page

REPORT



Staff prepare evaluation report

MPA managers contacted for outreach and contributions

Expert comment via Blue Parks web page

Evaluation and public comments forwarded to Science Council

DECISION




Science Council decides award status of nominees

AWARD



Nominees earning a Blue Park Award are recognized & celebrated by System partners







The Award Criteria

The Award Criteria









Blue Park Award Criteria A considerable body of scientific literature has focused on the attributes of protected areas that result in biodiversity conservation (e.g., Lester et al. 2009, Edgar et al. 2014, Zupan et al. 2018). International commitments, including the Convention on Biological Diversity, highlight the need for protected areas to be located in places important for biodiversity, effectively and equitably managed, ecologically representative and designed to support population connectivity and migration. Despite widespread interest in implementing effective MPAs, science has not typically informed the implementation of MPAs (Watson et al. 2016).

Blue Park standards rely on up-to-date science on the attributes of effective MPAs. There are three parts to the Blue Park evaluation. Nominees must meet the criteria in Part 1 to qualify for the award. The criteria in Part 2 determine the level of Blue Park Award that the nominee earns. The third part of the review process is an assessment of the conservation value the nominee adds to the network of Blue Parks via ecosystem representation and ecological spatial connectivity. Part 3 will prioritize nominees and new conservation targets that contribute most to safeguarding marine biodiversity.

Many of the criteria included in the Blue Park evaluation require a qualitative assessment of an MPA's attributes, as quantitative thresholds will depend on context. It is the role of the Science Council members – experts from across relevant disciplines and geopolitical boarders – to judge the degree to which each potential nominee meets the standards established by the Blue Park criteria. Criteria for which some circumstances are anticipated to affect the Science Council's judgement include a section entitled, "Additional Considerations."

ELIGIBILITY CRITERIA 1.1 Biodiversity Value 1.2 **Implementation** 0







Answers to 6 key questions determine the award status of a nominated site:

- 1.1 Does the MPA protect a place that is valuable for marine biodiversity?
- 1.2 Is the MPA designated for bio diversity conservation, and is it fully implemented?
- 2.1 Does the site prohibit activities that degrade the marine environment?
- 2.2 Is the design, governance and management of the MPA consistent with effectiveness, and are local communities represented in the management?
- 3.1 What value does the MPA add to the Blue Park network in terms of replicate ecosystem representation?
- 3.2 What value does the MPA add to the Blue Park network in terms of ecological spatial connectivity?

Eligibility Criteria



Blue Park Award Criteria



- http://www.iucnredlist.org/
- https://www.fws.gov/endangered/
- 3 http://www.ramsar.org/
- 4 https://maps.birdlife.org/ marineIBAs/default.html
- ⁵ https://www.mission-blue.org/ hope-spots/
- ⁶ http://www.cep.unep.org/ content/about-cep/spaw
- 7 http://www.natura.org/about.html

1.1 Biodiversity Value

Must satisfy at least one biodiversity value criterion

- 1.1.1 Includes rare, unique or representative ecosystems
- 1.1.2 Includes area of high species richness or endemism within the context of the biogeographic region
- 1.1.3 Includes demonstrated historic or predicted ecological (e.g., climate) refugia or populations with known or predicted tolerance or adaptive potential
- 1.1.4 Includes area important for threatened species (including those identified by the IUCN Red List¹ or national legislation), keystone species or foundational species these may be migration pathways or breeding, nursery, feeding or assembly areas.

Qualifying Designations

An MPA that has earned any of the following designations has satisfied at least one of the biodiversity value criteria above (1.1.1-1.1.4).

Convention on Wetlands of International Importance (Ramsar)³ Important Bird Area (IBA)⁴

Hope Spot⁵

Specially Protected Area and Wildlife (SPAW)⁶ in the Wider Caribbean Region Natura 2000 Special Protection Area (SPA) or Special Area of Conservation (SAC)⁷

1. Eligibility Criteria



Blue Park Award Criteria



Additional Consideration: Community-based MPAs

If a community-based MPA is longstanding and appears to be durable and permanent, the Science Council may consider it eligible for a Blue Park Award without government designation.

1 http://www.iucn.org/theme/ protected-areas/our-work/ green-list

1.2 Implementation:

Must satisfy all

- 1.2.1 The MPA is designated by a legitimate and functional government representing the interests of civil society, and the MPA's implementation meets the IUCN standards for recognizing indigenous peoples' rights.
- 1.2.2 The MPA is designated to enhance the biodiversity value of the site.
- 1.2.3 The MPA designation is permanent or is effective for at least 25 years.
- 1.2.4 The MPA has a management plan that has been updated within the last 15 years.
- 1.2.5 The MPA implements strategies to achieve high compliance that are appropriate to its ecological context, size and threats. These strategies must include clear communication to users and local communities about the regulations and MPA zone boundaries as well as enforcement strategies appropriate for the MPA's size, location, and poaching threats. Additional strategies may include incentivizing compliance or leveraging social influence through community engagement. Managers report high compliance in the MPA, and available evidence does not indicate high levels of non-compliance.
- 1.2.6 The MPA has a budget and staff.

Qualifying Designation

An MPA that has earned the following designation has satisfied all of the implementation criteria above (1.2.1-1.2.6).

IUCN Green List of Protected Areas¹

Ш

2. Award Status Criteria



Blue Park Award Criteria




Each eligible nominee will be evaluated in terms of its regulations and its design and management attributes. Note that a lower MPA Regulations score and a higher MPA Design, Management & Compliance score are needed to earn a more prestigious Blue Park Award. Platinum, Gold and Silver Awards require the following evaluation scores:

Blue Park Awards

Award Status	2.1 Regulations	2.2 Design, Management & Compliance	
Platinum	MPA regulation ≤3 score	AT LEAST 5 Design, Management & Compliance criteria met	
Gold	MPA regulation score ≤4	AT LEAST 4 Design, Management & Compliance criteria met	
Silver	MPA regulation score ≤5	AT LEAST 3 Design, Management & Compliance criteria met	

2.1 Regulations



Blue Park Award Criteria



Activities allowed in the MPA must be addressed by regulations in a management plan. Each regulatory zone of the MPA will be scored using the Regulation-Based Classification System for Marine Protected Areas (Costa et al. 2016), which relies on the number of fishing gear types allowed, a fishing gear impact score, bottom exploitation and aquaculture allowed, and anchoring and boating allowed. All zone scores are weighted according to their area and summed to generate an MPA score. Lower scores represent stronger levels of protection.

Additional Consideration : Other Activities

The MPA score serves as a guideline for the Science Council members in determining the award status of a nominee. Because the Regulation-Based Classification System for Marine Protected Areas is an estimate that sometimes conflates activities with very different impacts (e.g., salmon farming and oyster farming, SCUBA spearfishing and snorkel spearfishing) and excludes some damaging non-extractive recreational activities (e.g., snorkeling that results in high-impact coral reef trampling), the Science Council may decide a nominee's award status differently than the nominee's MPA score indicates.




Additional Consideration : Buffers

Buffer zones – zones of reduced human impact surrounding core no-take protected areas – enhance the conservation value of core no-take areas.

An MPA that includes a large (>100 km²) no-take zone (Zone regulation score 1-3) surrounded by a buffer zone with a score 3-5 may be considered for a Platinum Blue Park Award.

Blue Park Award Criteria





The following decision tree is used to determine the zone score for each zone of an MPA. Fishing gear impact scores, bottom exploitation & aquaculture index, and anchoring & boating index are determined according to the subsequent tables.

Zone Classification Number of 1-5 0 6-10 _ _ ≥ 15 20 20 fishing gears Fishing gear 6-8 0 ≤5 ≤8 impact score Bottom 0 $2 \leq 1 \leq 1 \leq 2 \leq 1 \leq 2$ ≤1 2 exploitation & aquaculture **Anchoring** 1 2 & boating **Zone Score** 3 4 5 1 2 4 6 6 6 5

Fishing Gear Impact Score

Use highest score among faishing gears allowed in the regulatory zone



Intertidal hand captures

Spearfishing/ diving



Fish traps

Fixed fish traps "modrogue"

Gillnets



Hand harvesting

Longlines (pelagic)

Traps (lobster/octopus/crab)



Dredges



Hand dredges (bivalves)

Lines (jigs, hook andline, rod, troll)

————
Purse seines

(pelagic)

Trawl (pelagic)

Longlines (bottom)



Beach seines

Surrounding nets near shore

Trammel net



Purse seines (bottom)

Trawl (bottom)


Bottom Exploitation & Aquaculture Index

Aquaculture and bottom exploitation not allowed	0
Aquaculture OR bottom exploitation allowed, but not mining/oil platforms/sand extraction/detonations	1
Both aquaculture AND bottom exploitation allowed, or only bottom exploitation including mining/oil platforms /sand extraction/detonations	2

Anchoring & Boating Index

No anchoring allowed	0
Boating and/or anchoring allowed, but anchoring is fully regulated: restricted to particular areas or mooring buoys	1
Boating and/or anchoring allowed, and anchoring is partially regulated or unregulated	

2.2 Design, Management & Capacity



Effective MPAs have a combination of design, management and compliance attributes. Nominees must have at least 5 of these attributes to be considered for a Platinum Award, 4 to be considered for a Gold Award and 3 to be considered for a Silver Award.



2.2.1 Size	MPA ≥ Explicitly designed as part of a network of MPAs to support population connectivity	
2.2.2 Ecological Isolation	Ecological or other protected area buffers surround habitats targeted for conservation within the MPA (e.g., soft sediment or deep water surrounding coral reefs).	
2.2.3 Age	The MPA is ≥ 10 years old.	
2.2.4 Effective Management Planning	The management plan identifies: 2.2.4.1 Measurable conservation targets 2.2.4.2 Threats to the conservation targets 2.2.4.3 Planned activities to mitigate threats and achieve conservation targets 2.2.4.4 Monitoring plans to measure progress towards conservation targets.	
2.2.5 Community Engagement	The local community is engaged in the management of the MPA.	
2.2.6 Resources & Capacity	The MPA has adequate resources and capacity (including budget, staff, training and leveraged partnerships) to implement its management plan and its compliance strategies.	

3. System Priorities



Blue Park Award Criteria



To ensure the representation of all ecosystems in all biogeographic regions, the replication of representative habitats, and the ability of the Blue Park network to support marine population persistence and migration, the Blue Park evaluation includes an assessment of each nominee's contribution to the conservation value of the network.

These geographic analyses will be conducted by program staff members using information about the existing Blue Parks. As the network grows, program staff will perform gap analyses to prioritize locations for new Blue Parks.

3.1 Ecosystem Representation

Blue Parks prioritizes nominees and Accelerator projects protecting ecosystems that are under-represented within their biogeographic region (< 30% are protected in Blue Parks), or are rare in the biogeographic region.

3.2 Ecological Spatial Connectivity

Blue Parks also prioritizes nominees and Accelerator projects that improve ecological spatial connectivity among existing Blue Parks. Ecological spatial connectivity refers to the physical and biological processes connecting areas in the marine environment in ways that support wildlife and ecosystems (Carr et al. 2017). System connectivity analyses are focused on population connectivity and migration; they rely on dispersal and migration estimates for key taxa as well as distances between Blue Parks with relevant habitats.

BLUE . PARKS



The Scientific Basis for Blue Park Award Criteria

The Scientific Basis for Blue Park Award Criteria





The Scientific Basis for Blue Park Award Criteria The Blue Park evaluation process is intended to identify MPAs that will help safeguard marine life. It is based on the best available science and has been developed over several years with the help of many marine scientists working in academia, government agencies, and non-profit conservation organizations. To ensure the Blue Park Award criteria continue to inform MPA implementation with the best available science, they will be updated regularly with the newest research. The following sections summarize the key scientific priciples of the Blue Park criteria.

Eligibility Criteria

1.1 Biodiversity Value

To make a significant contribution to protecting marine biodiversity, MPAs must protect biologically valuable places (Devillers et al. 2014). Many MPAs are currently located in places with sub-optimal biodiversity value because their protection involves fewer political hurdles, while highly biodiverse places or areas that support threatened or endangered species remain unprotected (Agardy et al. 2011, Devillers et al. 2014). The criteria for biodiversity value closely align with many other established criteria for MPA biodiversity value (Asaad et al. 2017), and therefore, an MPA that has earned one of the qualifying designations (see Criterion 1.1 in Section III) has already demonstrated it meets one or more of the Biodiversity Value criteria. For areas where there is a lack of information on the biodiversity value as it relates to regional or global patterns, additional research may be necessary. In addition to ecological refugia that may provide significant biodiversity conservation value in the context of current and future climate change, protecting populations with greater adaptive potential will promote long-term persistence and increase resilience to climate change (Walsworth et al. 2019, Selmoni et al. 2020).

The Scientific Basis for Blue Park Award Criteria

1.2 Implementation

MPAs can only become Blue Parks if they are fully implemented and designated for biodiversity conservation. Evidence of active management for biodiversity conservation includes stated objective to conserve or enhance biodiversity in the official documentation for the marine MPA (e.g., the designation legislation, the management plan or the official website). Additional evidence of the objective may include an updated management plan, a management team, a budget and an appropriate approach to enforcing protections. Blue Park Awards incentivize protections intended to be permanent that will result in long-term biodiversity conservation benefits.

To be eligible for a Blue Park Award, an MPA located in a place where indigenous people are present or have a collective attachment must comply with the IUCN Standard on Indigenous People (2006). This standard upholds indigenous peoples' rights to self-determination, the use of traditional ecological knowledge and consultation in management. This standard expressly prohibits the forced relocation of indigenous peoples for the purposes of conservation.

High compliance is essential to effective MPAs (Rife et al. 2013, Edgar et al. 2014, Bergseth et al. 2015), but challenging to measure. The Blue Park evaluation of compliance (1.2.5) relies on multiple indicators of compliance strategies and compliance outcomes - some qualitative and some quantitative - because each one has biases and provides an incomplete picture of compliance (Bergseth et al. 2015). In addition to assessing the degree to which compliance strategies (defined broadly, see Bergseth et al. 2017) are appropriate to the ecological context of the MPA as well as its threats, the Blue Parks evaluation includes a consideration of at least two years of enforcement records and enforcement effort (lacarella et al. 2021) as well as managers' self-assessments of enforcement effort and compliance rates. Self-assessment questions include the one used by Gill et al. (2017) as a measure of enforcement capacity and compliance because their analysis revealed a significant association between enforcement capacity and conservation outcomes - this question is taken from the Management Effectiveness Tracking Tool (Stolton et al. 2007) and the World Bank MPA Score Card (Stolton et al. 2007). The second and third self-assessment questions are based on those used by Bergseth et al. (in prep) to measure MPA compliance.

Self-assessment questions for managers:

- (1) How would you characterize the level of enforcement and compliance with regulations in the MPA? (Please select the statement that most closely resembles enforcement and compliance in the MPA.)
 - a) There are few or no rules or the staff lacks the capacity to enforce rules and regulations
 - b) There are rules and regulations, but they are inconsistently enforced
 - c) There are rules and regulations that are actively and consistently enforced
- (2) How many people fish illegally in no-take areas or using banned fishing gears in the marine protected area?
 - a) None
 - b) A small percent of the total users (less than 10%)
 - c) A large percent of the total users (10-50%)
 - d) Most of the total users (more than 50%)
- (3) How often do people fish illegally in no-take areas or using banned fishing gears in the marine protected area?
 - a) Never
 - b) Rarely (few occurrences)
 - c) Often
 - d) All the time

Where available, third-party expert assessment of compliance, scientific literature related to enforcement, additional compliance evidence collected by managers, and remote sensing data related to compliance will be considered in the Blue Park evaluation. The evaluation prioritizes ecological data showing a positive MPA effects (e.g., peer-reviewed research demonstrating higher fish abundance or biomass in the MPA), since these positive outcomes are associated with high compliance and reflect the

Additional Consideration

Community-based MPAs can be effective at conserving marine biodiversity (Chirico et al. 2017) and may be sustained through long-term community commitments rather than government designation. Therefore, the Science Council may consider a durable community-based MPA eligible for a Blue Park Award without government designation.

Award Status Criteria

2.1 Regulations

To effectively safeguard marine biodiversity, an MPA must regulate activities that negatively impact the biodiversity values of the site. The Blue Park evaluation prioritizes fully protected MPAs that exclude extractive activities (Lubchenco & Grorud-Colvert 2015), as these MPAs result in the best biological and ecological outcomes (Lester and Halpern 2008, Edgar et al. 2014, Appolloni et al. 2017, Strain et al. 2018, Aalto et al. 2019). Platinum Awards are reserved for these fully protected MPAs or those with a significant (> 100 km2) fully protected zone buffered by a zone allowing only very limited lower-impact extraction. However, 94% of all MPAs allow some fishing (Costello & Ballantine 2015), and the Blue Park criteria acknowledge the contributions of partially protected marine areas with strong regulations for biodiversity conservation (Lester & Halpern 2008, Coll et al. 2011, Tyler et al. 2011, Sciberras et al. 2013, Campbell et al. 2017, Gill et al. 2017, Giakoumi et al. 2017, Zupan et al. 2018), while recognizing that not all partially protected areas are effective at conserving fished populations (e.g., Denny and Babcock 2003, Di Franco et al. 2009). Gold and Silver awards are earned by highly protected areas (Horta e Costa et al. 2016, Zupan et al. 2018) designed, managed and enforced to contribute to biodiversity conservation.

The Scientific Basis for Blue Park Award Criteria

The Scientific Basis for Blue Park Award Criteria To assess the strength of an MPA's regulations, the Blue Park evaluation employs a classification system based on the number of fishing gears allowed, their ecological impact, the types of bottom exploitation and aquaculture allowed and the regulations relating to recreational boating (Horta e Costa et al. 2016, Zupan et al. 2018). For MPAs with multiple zones, the evaluation uses a weighted average of the individual zone scores (weighted by the proportion of zone area to total MPA area). The scores produced by this simple classification system strongly correlate with scores produced by a classification system employing a rigorous assessment of all MPA regulations (Horta e Costa et al. 2016) and also correlate with the biological outcomes of MPAs (Zupan et al. 2018).

Additional Consideration: Other Activities

Science Council members may consider allowed activities and regulated activities that are not well-represented by the regulations-based MPA classification system in determining the award status of a nominee. Of particular concern may be MPAs in which non-extractive activities have a large impact on the protected ecosystem (Thurston et al. 2012), such as coral reef trampling by tourists (Williamson et al. 2016).

Additional Consideration: Buffers

Buffer zones – areas with some fishing restrictions surrounding no-take zones – do not necessarily contribute positively to the conservation value of an MPA (Claudet et al. 2008, Di Franco et al. 2009). However, a strongly regulated zone – one that would earn a Blue Park Award on its own – buffering a very large fully protected zone (> 100 km²) may be considered for a Platinum Blue Park Award, given that strictly regulated buffer zones can effectively enhance conservation outcomes (Coll et al. 2011).

2.2 Design, Management, & Capacity

MPA design, management and compliance play important roles in MPA effectiveness (e.g., Claudet et al. 2008, Le Quesne 2009, Gaines et al. 2010, McCook et al. 2010, Edgar et al. 2014). Attributes that are associated with positive biological and ecological outcomes should be standard practice in MPA implementation. To this end, the Blue Park standard requires at least 3 of the 6 attributes outlined below for Silver status, 4 for Gold status and 5 for Platinum status.

2.2.1 Size

Small no-take MPAs can have positive conservation effects (e.g., Giakoumi et al. 2017), however larger MPAs are better because they support larger population sizes (Gaines et al. 2010), they are more likely to protect self-persistent populations sustained by the local retention of larvae (Botsford et al. 2003), they are more likely to exceed the home range size of fished species (Di Franco et al. 2018), and they are more likely to contain fully functional ecosystems (McLeod et al. 2009). Larger MPAs have stronger positive impacts on biological communities (Claudet et al. 2008, Friedlander et al. 2017), particularly when those MPAs exhibit two or more other attributes of effective MPAs effective management, longevity, isolation and no-take restrictions (Edgar et al. 2014). Using individual-based models to evaluate effectiveness of different sizes of no-take MPAs in coastal coral reef ecosystems, Krueck et al. (2017) determined that an MPA needed to be at least 2-10 km wide to achieve partial protection of fished species and 100 km wide to achieve full protection of nearly all the species they modeled. Claudet et al. (2008) and Edgar et al. (2014) defined large MPAs as > 100 km² in their analyses of MPA effectiveness. We will consider nominees > 100 km² for consideration of higher status Blue Park Awards than those < 100 km2 until a more nuanced assessment of the relationship between size and conservation effectiveness across a variety of ecosystems is published.



An effective alternative to a large MPA is a network of smaller MPAs spaced closely enough to support population connectivity for species with longer dispersal distances (Gaines et al. 2010, McCook et al. 2010, Grorud-Colvert et al. 2014, Carr et al. 2017, Baetscher et al. 2019) and sized large enough to support the local retention of species with shorter dispersal distances (Carr et al. 2017).

2.2.2 Ecological Isolation

Ecosystems entirely enclosed within protected area boundaries and isolated by deep water, sand or other ecological barriers provide more conservation benefits (Edgar et al. 2014). We will also consider ecological barriers in adjacent protected areas (e.g., terrestrial protected areas).

2.2.3 Age

Older MPAs – those over 10 years old (Claudet et al. 2008, Edgar et al. 2014, Strain et al. 2018) or 15 years old (Molloy et al. 2009) – confer greater conservation benefits than younger MPAs (Friedlander et al. 2017). Depending on local environmental and species-specific population drivers, some species may not show a measurable response to protection for at least a decade (Kaplan et al. 2019). An MPA younger than 10 years old that earns Blue Park status may be eligible to earn a higher award status

2.2.4 Management Planning

Employing a management plan has been associated with positive ecological outcomes in tropical MPAs (Hargreaves-Allen et al. 2017). Effective management planning and governance involves identifying and communicating measurable objectives, prioritized threats and management activities, including monitoring for ecological outcomes (Tear et al. 2005, Bennett and Satterfield 2018). Blue Parks must design and implement an ecological monitoring program that measures progress with re-

2.2.5 Community Engagement

Though more research is needed to determine the relationship between participatory governance and ecological outcomes in MPAs (Stafford 2018), community consultation in the implementation and management of an MPA is associated with higher levels of compliance with MPA regulations (Pollnac et al. 2010, Eriksson et al. 2019) and MPA effectiveness (Batista and Cabral 2016).

2.2.6 Resources & Capacity

The capacity to carry out all management and compliance activities is vital to the conservation performance of MPAs. Adequate staff capacity and budget is particularly important for MPA effectiveness (Gill et al. 2017, Bennett and Satterfield 2018).

System Priorities

The conservation value of an MPA depends, in part, on its spatial relationship to other MPAs (McCook et al. 2009, McLeod et al. 2009, Gaines et al. 2010, Batista and Cabral 2016). Blue Park criteria incorporate a geographic framework with an assessment of ecosystem representation and ecological connectivity to enhance biodiversity persistence (Magris et al. 2018), with a goal of 30% protection for each habitat type within each biogeographic region. Assessing the network value of a nominee is different from the other parts of the evaluation because it depends on an analysis of the existing protections for the ecosystems within the nominee's biogeographic region.

The Scientific Basis for Blue Park Award Criteria

A biogeographic region is a large area defined by biotic distributions. Marine Conservation Institute is building both a network and a portfolio of well-protected Blue Parks distributed across the coastal biogeographic regions defined by Spalding et al. (2007). Marine Conservation Institute is also working with partners to develop a more comprehensive three-dimensional ecoregional framework for the global ocean (Sayre et al. 2017).

3.1 Replicate Ecosystem Representation


The Blue Park network must include examples of all the marine ecosystems in each biogeographic region of the ocean to protect the vast diversity of marine life (Spalding et al. 2008, McCook et al. 2009, McLeod et al. 2009), because within each region, places with comparable physical and chemical conditions tend to host similar species assemblages. The Blue Park network will include replicate sites of each ecosystem within a biogeographic region to protect against unforeseen ecological disasters and guard against biodiversity loss (Gaines et al. 2010). The goal is to include requisite sites to achieve the conservation target of at least 30% of each ecosystem within each biogeographic region, and up to 100% of particularly rare and threatened ecosystems (O'Leary et al. 2016). As the Blue Park network grows, program staff will perform gap analyses to identify priority ecosystems within each biogeographic region to target in new MPAs.

The Scientific Basis for Blue Park Award Criteria

3.2 Ecological Spatial Connectivity

Marine Conservation Institute aims to assemble a network of effective Blue Parks that support marine population connectivity and migration. Avoiding extinction requires either a population's continuous presence in a habitat or the ability to recolonize after local extinction. Population persistence in the ocean often depends on network persistence among demographically connected populations (Botsford et al. 2001, Hastings and Botsford 2006, McCook et al. 2009, McCook et al. 2010). Networks of MPAs add resilience for protected populations that are demographically connected (Moffitt et al. 2011) and for ecosystems that are ecologically connected (Carr et al. 2017). MPA networks can also provide opportunities for adaptive migration in response to climate stressors (Hole et al. 2011, Fredston-Hermann et al. 2018), and conserving sources of recolonizers is important when pulse stressors (e.g., acute coral bleaching events, storms or upwelled low-oxygen waters) cause local extinctions (Gaines et al. 2003) or mass mortality of density-sensitive species (Aalto et al. 2019). Therefore, protecting enough cumulative area and enough dispersed replicates of ecosystems that could be connected via larval, sub-adult or adult movement is an effective means of building portfolio resilience into the network of Blue Parks (McCook et al. 2009, Grorud-Colvert et al. 2014, Carr et al. 2017, Baetscher et al. 2019).

BLUE . PARKS



Acknowledgements

Blue Parks Partners

The Blue Parks program relies on the contributions of many diverse partners. The Blue Park Award criteria, in particular, depend on the contributions of many scientists and MPA experts from across disciplines and sectors. These criteria were first developed based on the scientific literature and on expert opinion elicited in four focused workshops hosted by Marine Conservation Institute over three years. Program staff and Science Council members review and update these criteria annually. Marine Conservation Institute is very grateful for the important contributions made by many scientists and conservation experts around the world.

BLUE . PARKS



References

VI

Aalto, E. A., F. Micheli, C. A. Boch, J. A. E. Montes, C. B. Woodson, and G. A. De Leo. 2019. Catastrophic mortality, allele effects, and marine protected areas. The American Naturalist 193:391–408.

Agardy, T., G. N. di Sciara, and P. Christie. 2011. Mind the gap: Addressing the shortcomings of marine protected areas through large scale marine spatial planning. Marine Policy 35:226–232.

Appolloni, L., S. Bevilacqua, L. Sbrescia, R. Sandulli, A. Terlizzi, and G. F. Russo. 2017. Does full protection count for the maintenance of ß-diversity patterns in marine communities? Evidence from Mediterranean fish assemblages. Aquatic Conservation: Marine and Freshwater Ecosystems 27:828–838.

Asaad, I., C. J. Lundquist, M. V. Erdmann, and M. J. Costello. 2017. Ecological criteria to identify areas for biodiversity conservation. Biological Conservation 213:309–316.

Baetscher, D. S., E. C. Anderson, E. A. Gilbert-Horvath, D. P. Malone, E. T. Saarman, M. H. Carr, and J. C. Garza. 2019. Dispersal of a nearshore marine fish connects marine reserves and adjacent fished areas along an open coast. Molecular Ecology 28:1611-1623.

Ban, N. C., G. G. Gurney, N. A. Marshall, C. K. Whitney, M. Mills, S. Gelcich, N. J. Bennett, M. C. Meehan, C. Butler, S. Ban, T. C. Tran, M. E. Cox, and S. J. Breslow. 2019. Well-being outcomes of marine protected areas. Nature Sustainability 2:524-532.

Barr, L. M., R. L. Pressey, R. A. Fuller, D. B. Segan, E. McDonald-Madden, and H. P. Possingham. 2011. A New Way to Measure the World's Protected Area Coverage. PLoS ONE 6:e24707.

Batista, M. I., and H. N. Cabral. 2016. An overview of Marine Protected Areas in SW Europe: Factors contributing to their management effectiveness. Ocean Coastal Management 132:15–23.

Bennett, N. J. and T. Satterfield. 2018. Environmental governance: A practical framework to guide design, evaluation, and analysis. Conservation Letters 2018:e12600.

Bergseth, B. J. 2018. Effective marine protected areas require a sea change in compliance management. ICES Journal of Marine Science, 75:1178-1180.

Bergseth, B. J., G. R. Russ, and J. E. Cinner. 2015. Measuring and monitoring compliance in no-take marine reserves. Fish and Fisheries 16:240-258.

Di Franco, A., S. Bussotti, A. Navone, P. Panzalis, and P. Guidetti. 2009. Evaluating effects of total and partial restrictions to fishing on Mediterranean rocky-reef fish assemblages. Marine Ecology Progress Series 387:275–285.

Botsford, L. W., A. Hastings, and S.D. Gaines. 2001. Dependence of sustainability on the configuration of marine reserves and larval dispersal distance. Ecology Letters 4:144–150.

Botsford, L. W., F. Micheli, and A. Hastings. 2003. Principles for the design of marine reserves. Ecological Applications 13:S25–S31.

Campbell, S. J., G. J. Edgar, R. D. Stuart-Smith, G. Soler, and A. E. Bates. 2018. Fishing-gear restrictions and biomass gains for coral reef fishes in marine protected areas. Conservation Biology 32:401–410

Carr, M. H., S. P. Robinson, C. Wahle, G. Davis, S. Kroll, S. Murray, E. J. Schumacker, and M. Williams. 2017. The central importance of ecological spatial connectivity to effective coastal marine protected areas and to meeting the challenges of climate change in the marine environment. Aquatic Conservation: Marine and Freshwater Ecosystems 27:6–29.

Chirico, A. A. D., T. R. McClanahan, and J. S. Eklöf. 2017. Community- and government-managed marine protected areas increase fish size, biomass and potential value. PLoS ONE 12:e0182342.

Claudet, J., C. W. Osenberg, L. Benedetti-Cecchi, P. Domenici, J.-A. García-Charton, Á. Pérez-Ruzafa, F. Badalamenti, J. Bayle-Sempere, A. Brito, F. Bulleri, J.M. Culioli, M. Dimech, J. M. Falcón, I. Guala, M. Milazzo, J. Sánchez-Meca, P. J. Somerfield, B. Stobart, F. Vandeperre, C. Valle, and S. Planes. 2008. Marine reserves: size and age do matter. Ecology Letters 11:481–489.

Coll, J., A. Garcia-Rubies, G. Morey, and A. M. Grau. 2011. The carrying capacity and the effects of protection level in three marine protected areas in the Balearic Islands (NW Mediterranean). Scientia Marina 76:809–826.

Costello, M. J. and B. Ballantine. 2015. Biodiversity conservation should focus on no-take Marine Reserves. Trends in Ecology & Evolution 30:507–509.

Denny, C. M. and R. C. Babcock. 2004. Do partial marine reserves protect reef fish assemblages? Biological Conservation 116:119-129.

Di Franco, A., S. Bussotti, A. Navone, P. Panzalis, and P. Guidetti. 2009. Evaluating effects of total and partial restrictions to fishing on Mediterranean rocky-reef fish assemblages. Marine Ecology Progress Series 387:275-285.

Di Franco, A., J. G. Plass-Johnson, M. Di Lorenzo, B. Meola, J. Claudet, S. D. Gaines, J. A. García-Charton, S. Giakoumi, K. Grorud-Colvert, C. W. Hackradt, F. Micheli, and P. Guidetti. 2018. Linking home ranges to protected area size: The case study of the Mediterranean Sea. Biological Conservation 221:175-181.

Edgar, G. J., R. D. Stuart-Smith, T. J. Willis, S. Kininmonth, S. C. Baker, S. Banks, N. S. Barret, M. A. Becerro, A. T. F. Bernard, J. Berkhout, C. D. Buxton, S. J. Campbell, A. T. Cooper, M. Davey, S. C. Edgar, G. Försterra, D. E. Galván, A. J. Irigoyen, D. J. Kushner, R. Moura, P. E. Parnell, N. T. Shears, G. Soler, E. M. A. Strain, and R. J. Thomson. 2014. Global conservation outcomes depend on marine protected areas with five key features. Nature 506:216-220.

Eriksson, B., F. Johansson, and M. Blicharska. 2019. Socio-economic impacts of marine conservation efforts in three Indonesian fishing communities. Marine Policy 103:59-67.

Fernandez, L., J. Day, A. Lewis, S. Slegers, B. Kerrigan, D. Breen, D. Cameron, B. Jago, J. Hall, D. Lowe, and J. Innes. 2005. Establishing representative no-take areas in the Great Barrier Reef: largescale implementation of theory on marine protected areas. Conservation Biology 19:1733-44.

Fredston-Hermann, A., S. D. Gaines, and B. S. Halpern. 2018. Biogeographic constraints to marine conservation in a changing climate. Annals of the New York Academy of Sciences 367:49-13.

Friedlander, A. M., Y. Golbuu, E. Ballesteros, J. E. Caselle, M. Gouezo, D. Olsudong, & E. Sala. 2017. Size, age, and habitat determine effectiveness of Palau's Marine Protected Areas. PLoS ONE 12:e0174787.

Gaines, S. D., B. Gaylord, and J. L. Largier. 2003Avoiding current oversights in marine reserve design. Ecological Applications 13:32-46.

Gaines, S. D., C. White, M. H. Carr, and S. R. Palumbi. 2010. Designing marine reserve networks for both conservation and fisheries management. Proceedings of the National Academy of Sciences of the United States of America 107:18286-18293.

Gedan, K. B., M. L. Kirwan, E. Wolanski, E. B. Barbier, and B. R. Silliman. 2010. The present and future role of coastal wetland vegetation in protecting shorelines: answering recent challenges to the paradigm. Climatic Change 106:7-29.

Giakoumi, S., C. Scianna, J. Plass-Johnson, F. Micheli, K. Grorud-Colvert, P. Thiriet, J. Claudet, G. Di Carlo, A. Di Franco, S. D. Gaines, and J. A. García-Charton. 2017. Ecological effects of full and partial protection in the crowded Mediterranean Sea: a regional meta-analysis. Scientific Reports 7:8940.

Gill, D. A., M. B. Mascia, G. N. Ahmadia, L. Glew, S. E. Lester, M. Barnes, I. Craigie, E. S. Darling, C. M. Free, J. Geldmann, S. Holst, O. P. Jensen, A. T. Whote, X. Basurto, L. Coad, R. D. Gates, G. Guannel, P. J. Mumby, H. Thomas, S. Whitmee, S. Woodley, and H. E. Fox. 2017. Capacity shortfall hinder the perfomance of marine protected areas globally. Nature 543:665-669.

Grorud-Colvert, K., J. Claudet, B. N. Tissot, J. E. Caselle, M. H. Carr, J. C. Day, A. M. Friedlander, S. E. Lester, T. L. de Loma, D. Malone, and W. J. Walsh. 2014. Marine Protected Area Networks: Assessing Whether the Whole Is Greater than the Sum of Its Parts. PLoS ONE 9:e102298.

Guidetti, P., M. Milazzo, S. Bussotti, A. Molinari, M. Murenu, A. Pais, N. Spanò, R. Balzano, T. Agardy, F. Boero, G. Carrada, R. Cattaneo-Vietti, A. Cau, R. Chemello, S. Greco, A. Manganaro, G. Notarbartolo di Sciara, G. F. Russo, and L. Tunesi. 2008. Italian marine reserve effectiveness: Does enforcement matter? Biological Conservation 141:699-709.

Hargreaves-Allen, V. A., Mourato, S., and Milner-Gulland, E. J. 2017. Drivers of coral reef marine protected area performance. PLoS ONE 12:e0179394.

Hastings, A., and L. W. Botsford. 2006. Persistence of spatial populations depends on returning home. Proceedings of the National Academy of Sciences of the United States of America

103:6067-6072.

Hole, D. G., B. Huntley, J. Arinaitwe, S. H. Butchart, Y. C. Collingham, L. D. C. Fishpool, D. J. Pain, and

S. G. Willis. 2011. Toward a management framework for networks of protected areas in the face of

climate change. Conservation Biology 25:305-315.

VI

Horta e Costa, B., J. Claudet, G. Franco, K. Erzini, A. Caro, and E. J. Gonçalves. 2016. A regulation-based classification system for Marine Protected Areas (MPAs). Marine Policy 72:192–198.

International Union for the Conservation of Nature (IUCN). 2016. IUCN Standard on Indigenous Peoples, Version 2.0. IUCN Environmental and Social Management System (ESMS).

lacarella, J. C., G. Clyde, B. J. Bergseth, and N. C. Ban. 2021. A synthesis of the prevalence and drivers of non-compliance in marine protected areas. Biological Conservation 255:108992.

Kaplan, K.A., L. Yamane, L.W. Botsford, M.L. Baskett, A. Hastings, S. Worden, and J.W. White. 2019. Setting expected timelines of fished population recovery for the adaptive management of a marine protected area network. Ecological Applications 29:e01949.

Krueck, N. C., C. Legrand, G. N. Ahmadia, Estradivari, A. Green, G. P. Jones, C. Riginos, E. A. Treml, and P. J. Mumby. 2017. Reserve Sizes Needed to Protect Coral Reef Fishes. Conservation Letters 11:e12415.

Lester, S. E., and B. S. Halpern. 2008. Biological responses in marine no-take reserves versus partially protected areas. Marine Ecology Progress Series 367:49–56.

Lester, S. E., B. S. Halpern, K. Grorud-Colvert, J. Lubchenco, B. I. Ruttenberg, S.D. Gaines, S. Airame, and R. R. Warner. 2009. Biological effects within no-take marine reserves: a global synthesis. Marine Ecology Progress Series 384:33–46.

Le Quesne, W. J. F. 2009. Are flawed MPAs any good or just a new way of making old mistakes? ICES Journal of Marine Science 66:132–136.

Lubchenco, J., and K. Grorud-Colvert. 2015. Making waves: The science and politics of ocean protection. Science 350:382–383.

McCook, L. J., G. R. Almany, M. L. Berumen, J. C. Day, A. L. Green, G. P. Jones, J. M. Leis, S. Planes, G. R. Russ, P. F. Sale, and S. R. Thorrold. 2009. Management under uncertainty: guide-lines for incorporating connectivity into the protection of coral reefs. Coral Reefs 28:353–366.

Magris, R. A., M. Andrello, R. L. Pressey, D. Mouillot, A. Dalongeville, M. N. Jacobi, and S. Manel. 2018. Biologically representative and well-connected marine reserves enhance biodiversity persistence in conservation planning. Conservation Letters 96:e12439.

McCook, L. J., T. Ayling, M. Cappo, J. H. Choat, R. D. Evans, D. M. De Freitas, M. Heupel, T. P. Hughes, G. P. Jones, B. Mapstone, H. Marsh, M. Mills, F. J. Molloy, C. R. Pitcher, R. L. Pressey, G. R. Russ, S. Sutton, H. Sweatman, R. Tobin, D. R. Wachenfeld, and D. H. Williamson. 2010. Adaptive management of the Great Barrier Reef: a globally significant demonstration of the benefits of networks of marine reserves. Proceedings of the National Academy of Sciences of the United States of America 107:18278–18285.

McLeod, E., R. Salm, A. Green, and J. Almany. 2009. Designing marine protected area networks to address the impacts of climate change. Frontiers in Ecology and the Environment 7:362–370.

Mellin, C., M. A. MacNeil, A. J. Cheal, M. J. Emslie, and M. J. Caley. 2016. Marine protected areas increase resilience among coral reef communities. Ecology Letters 19:629–637.

Micheli, F., A. Saenz-Arroyo, A. Greenley, L. Vazquez, J. A. E. Montes, M. Rossetto, and G. A. De Leo. 2012. Evidence That Marine Reserves Enhance Resilience to Climatic Impacts. PLoS ONE 7:e40832.

Mizrahi, M., A. Diedrich, R. Weeks, and R. L. Pressey. 2019. A Systematic Review of the Socioeconomic Factors that Influence How Marine Protected Areas Impact on Ecosystems and Livelihoods. Society and Natural Resources 32:4-20.

Moffitt, E. A., J. W. White, and L. W. Botsford. 2011. The utility and limitations of size and spacing guidelines for designing marine protected area (MPA) networks. Biological Conservation 144:306–318.

Molloy, P. P., I. B. McLean, and I. M. Côté. 2009. Effects of marine reserve age on fish populations: a global meta-analysis. Journal of Applied Ecology 46:743–751.

Nagelkerken, I. and S. D. Connell. 2015. Global alteration of ocean ecosystem functioning due to increasing human CO2 emissions. Proceedings of the National Academy of Sciences of the United States of America 112:13272-13277.

VI

Pollnac, R., P. Christie, J. E. Cinner, T. Dalton, T. M. Daw, G. E. Forrester, N. A. J. Graham, and T. R. McClanahan. 2010. Marine reserves as linked social-ecological systems. Proceedings of the National Academy of Sciences of the United States of America 107:18262–18265.

Rife, A. N., B. Erisman, A. Sanchez, and O. Aburto-Oropeza. 2013. When good intentions are not enough... Insights on networks of "paper park" marine protected areas. Conservation Letters, 6:200-212.

Sayre, R. G., D. J. Wright, S. P. Breyer, K. A. Butler, K. Van Graafeiland, M. J. Costello, P. T. Harris, K. L. Goodin, J. M. Guinotte, Z. Basher, M. T. Kavanaugh, P. N. Halpin, M. E. Monaco, N. Cressie, P. Aniello, C. E. Frye, and D. Stephens. 2017. A three-dimensional mapping of the ocean based on environmental data. Oceanography 30:90–103.

Sciberras, M., S. R. Jenkins, R. Mant, M. J. Kaiser, S. J. Hawkins, and A. S. Pullin. 2013. Evaluating the relative conservation value of fully and partially protected marine areas. Fish and Fisheries 16:58–77.

Selmoni, O., G. Lecellier, L. Vigliola, V. Berteaux-Lecellier, and S. Joost. 2020. Coral cover surveys corroborate prediction on reef adaptive potential to thermal stress. Scientific Reports 10:19680.

Spalding, M. D., L. Fish, and L. J. Wood. 2008. Toward representative protection of the world's coasts and oceans-progress, gaps, and opportunities. Conservation Letters 1:217–226.

Spalding, M. D., H. E. Fox, G. R. Allen, N. Davidson, Z. A. Ferdana, M. Finlayson, B. S. Halpern, M. A. Jorge, A. L. Lombana, S. A. Lourie, K. D. Martin, E. McManus, J. Molnar, C. A. Recchia, and J. Robertson. 2007. Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. BioScience 57:573–583.

Stafford, R. 2018. Lack of evidence that governance structures provide real ecological benefits in marine protected areas. Oceanography 152:57–61.

Staub, F. and Hatziolos, M. E. 2004. Score Card to Assess Progress in Achieving Management Effectiveness Goals for Marine Protected Areas. World Bank.

Stolton, S., M. Hockings, N. Dudley K. MacKinnon, T. Whitten, and F. Leverington. 2007. Management Effectiveness Tracking Tool: Reporting Progress in Protected Area Sites; Second Edition. Gland, Switzerland: WWF International.

Strain, E. M. A., G. J. Edgar, D. Ceccarelli, R. D. Stuart-Smith, G. R. Hosack, and R. J. Thomson. 2018. A global assessment of the direct and indirect benefits of marine protected areas for coral reef conservation. Diversity and Distributions 10:1–12.

Tear, T. H., P. Karieva, P. L. Angermeier, P. Comer, B. Czech, R. Kautz, L. Landon, D. Mehlman, K. Murphy, M. Ruckelshaus, J. M. Scott, and G. Wilhere. 2005. How much is enough? The recurrent problem of setting measurable objectives in conservation. BioScience 55:835–849.

Thresher, R. E., J. M. Guinotte, R. J. Matear, and A. J. Hobday. 2015. Options for managing impacts of climate change on a deep-sea community. Nature Climate Change 5:635–639.

Thurstan, R. H., J. P. Hawkings, L. Neves, and C. M. Roberts. 2012. Are marine reserves and non-consumptive activities compatible? A global analysis of marine reserve regulations. Marine Policy 36:1096–1104.

Tyler, E. H. M., A. Manica, N. Jiddawi, and M. R. Speight. 2011. A role for partially protected areas on coral reefs: maintaining fish diversity? Aquatic Conservation: Marine and Freshwater Ecosystems 21:231–238.

Walsworth, T. E., D. E. Schindler, M. A. Colton, M. S. Webster, S. R. Palumbi, P. J. Mumby, T. E. Essington, and M. L. Pinsky. 2019. Management for network diversity speeds evolutionary adaptation to climate change. Nature Climate Change 9:632–636.

Watson, J. E. M., E. S. Darling, O. Venter, M. Maron, J. Walston, H. P. Possingham, N. Dudley, M. Hockings, M. Barnes, and T. M. Brooks. 2016. Bolder science needed now for protected areas. Conservation Biology 30:243–248.

Williamson, J. E., E. E. Byrnes, J. A. Clark, D. M. Connolly, S. E. Schiller, J. A. Thompson, L. Tosetto, J. C. Martinelli, and V. Raoult. 2016. Ecological impacts and management implications of reef walking on a tropical reef flat community. Marine Pollution Bulletin 114:742–750.

Credits

Photos

- ¹ Nicklen, Paul. *Polar bear.* 2016.
- ² Platko, Joe. 2016.
- ³ Platko, Joe. *Apo Turtle*. 2016.
- ⁴ Platko, Joe. *Hawai'i Reef Scene.* 2014.
- ⁵ Bauder, Clinton. Mono Lobo Wall: Monterey Scenics. 2016.
- ⁶ Nicklen, Paul. Seal. 2016.

Cover & Document Design

Chathurika Thenuwara